

IF YOU RECEIVED THIS FAX IN ERROR,
PLEASE CALL 604-806-8886 IMMEDIATELY



PRESCRIBER'S ORDERS

NO DRUG WILL BE DISPENSED OR ADMINISTERED
WITHOUT A COMPLETED
CAUTION SHEET
ALLERGY/INTOLERANCE STATUS FORM (PHC-PH047)

DATE
AND TIME

OPHTHALMOLOGY OPERATIVE ORDERS

[see corresponding Medication Administration Record PHC-PH270-MA (R. Feb 28-17)]
(Items with check boxes must be selected to be ordered)

OPERATIVE EYE: Left Right

PRE-OPERATIVE

Monitoring: Temp, BP, Pulse, RR, O₂ Saturation. Monitoring as per DT Procedure Rooms, Patient Admission and Pre-operative Assessment protocol

Capillary Blood Glucose on admission (Diabetic patient)

MEDICATIONS:

Anxiety: lorazepam 0.5 mg *OR* 1 mg sublingually on arrival

lorazepam 0.5 to 1 mg sublingually PRN once for moderate to high anxiety

Ophthalmic:

tetracaine 0.5% ophthalmic solution x 1 drop into operative eye once

cyclopentolate 1% ophthalmic solution x 2 drops into operative eye; repeat dose after 5 to 15 minutes

moxifloxacin 0.5% ophthalmic solution x 2 drops into operative eye; repeat dose after 5 to 15 minutes

PHENYLEphrine 2.5% ophthalmic solution x 2 drops into operative eye; repeat dose after 5 to 15 minutes

If slow dilation: PHENYLEphrine 10% ophthalmic solution x 1 drop into operative eye once

atropine 1% ophthalmic solution x 2 drops into operative eye; repeat dose after 5 to 15 minutes

pilocarpine 1% ophthalmic solution x 1 drop into operative eye; repeat dose after 5 to 15 minutes

lidocaine 2% jelly to be applied to operative eye by surgeon

PROCEDURAL SEDATION:

Saline lock

fentanyl 25 mcg IV PRN to maximum of 50 mcg

midazolam 0.5 mg IV PRN to maximum of 1 mg

Monitor patient as per Procedural Sedation in Clinics and Procedure Rooms, protocol

Discharge patients when procedural sedation discharge criteria met per Procedural Sedation in Clinics and Procedure Rooms, protocol

RETRO OR PERIBULBAR BLOCKS: (administered by ophthalmologist)

Saline lock

lidocaine 2% with epinephrine 1:100,000 2 to 4 mL *AND*

bupivacaine 0.75% 2 to 4 mL;

may repeat to maximum of 8 mL combined total

POST-OPERATIVE:

Monitoring: Monitor patient as per DT Procedure Rooms, Post op care discharge scoring assessment

acetazolamide 250 to 500 mg PO once immediately post-op

acetaminophen 325 to 650 mg PO Q3 to 4H PRN pain

TYLENOL #3 1 to 2 tablets PO Q4H PRN pain

patient must remain supine x _____ minutes post-operatively

Printed Name

Signature

College ID

Contact Number



ALL NEW ORDERS MUST BE FLAGGED